

HEDIS Hints

Appropriate Testing for Pharyngitis (CWP)

Appropriate Testing for Pharyngitis (CWP)

- What does NCQA's Appropriate Testing for Pharyngitis (CWP) Measure?
- Examples of Best Practices Documentation
- Coding to Improve this HEDIS Measure
- Questions

What is NCQA's CWP Measure?

- The CWP measure monitors the percentage of episodes for members 3 years and older where the member was:
 - diagnosed with pharyngitis
 - dispensed an antibiotic
 - and received a group A streptococcus (strep) test
- The look back period for this measure begins July 1 of the year prior to the measurement year and ends on June 30 of the measurement year.
- This measure looks for the date of service for any outpatient, telephone, observation or ED visit, e-visit or virtual check-in during the look back period with a diagnosis of pharyngitis.

What is NCQA's CWP Measure?

- The age stratifications for this measures are:
 - 3-17 years old
 - 18-64 years old
 - 65 years and older
 - Total (sum of all stratifications)
- A member may be in this measure for multiple episodes.
- This measure **is only an administrative measure** which means that the only way to get credit for this is to submit the appropriate code on a claim.
- Health Partners Plans **may not perform** medical record review for the AAP measure.

Examples of Best Practices Documentation

- Perform a rapid strep test and/or a throat culture to confirm diagnosis before prescribing antibiotics.
- Do not treat any sign/symptoms of possible strep throat with antibiotics unless a rapid strep test and/or a throat culture is performed.
- Document results of the rapid strep test and/or they throat culture.
- Document refusals of testing for strep by members.
- Submit co-morbid diagnosis coded that apply to the visit on the claim.
- Document and code accurately if antibiotics are prescribed for another condition.

Examples of Best Practices Documentation

- Educate patients, parents and families on:
 - Difference between viral and bacterial infections.
 - Inappropriate use of antibiotics.
 - Explain other causes for sore throats such as colds or allergies.
 - Offers ways to treat symptoms such as:
 - Increased fluids by mouth
 - Over the counter medicines
 - Cool mist vaporizer
 - Throat lozenges
 - Rest
 - Explain ways to prevent infection:
 - Wash hands frequently
 - Clean children's toys
 - Wipe down frequently touched areas with disinfectant wipes

Coding to Improve this HEDIS Measure

- **ICD-10:**
 - Pharyngitis: J02.0, J02.8, J02.9, J03.00, J03.01, J03.80, J03.81, J03.90, J03.91
- **CPT:**
 - Group A strep tests: 87070, 87071, 87081, 87430, 87650, 87651, 87652, 87880
 - Online assessments: 98970, 98971, 98972, 99421, 99422, 99423, 99457
 - Telephone visits: 98966, 98967, 98968, 99441, 99442, 99443
- **HCPCS:**
 - Online assessments: G0071, G2010, G2012

Questions?

Please contact ProviderEducation@hpplans.com

or

Teresa McKeever MS, BSN, RN
Director, Quality Management
tmckeever@hpplans.com